

## Original Research Article

# Lens thickness and associated ocular biometric factors among cataract patients in tertiary hospital

Shivani Sharma, Anil Kumar Sharma, Satish Gupta, Divyanshi Gupta\*

Department of Ophthalmology, GMC Jammu, Jammu and Kashmir, India

**Received:** 15 January 2023

**Revised:** 29 January 2023

**Accepted:** 30 January 2023

**\*Correspondence:**

Dr. Divyanshi Gupta,

E-mail: Divyanshi.gupta95@gmail.com

**Copyright:** © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

### ABSTRACT

**Background:** A significant factor in determining intraocular lens power is the biometry assessment of cataract patients prior to surgery. To evaluate the distribution of lens thickness (LT) and its associations with other ocular biometric factors among cataract patients.

**Methods:** Total 978 eyes from cataract patients were retrospectively included. Ocular biometric factors including k1, k2, LT, central corneal thickness (CCT), anterior chamber depth (ACD), white-to-white (WTW) distance, and axial length (AL) were collected based on the medical records. The associations between LT and general or ocular factors were assessed. We analyzed the data using descriptive analysis and correlated each variable using the Spearman's Rho analysis.

**Results:** The mean age was  $62.5 \pm 15.70$  years and 55.11% were females. Mean LT was  $4.35 \pm 0.41$  mm. The LT was greater in older patients. LT positively correlated with AL, WTW and CCT, while negatively correlated with ACD.

**Conclusions:** LT is an important biometric parameter that should be considered, along with other biometric parameters, in determining effective lens position in patients undergoing cataract surgery. ACD is predominantly influenced by LT than the AL.

**Keywords:** ACD, AL, Biometry, Intraocular lens, LT

### INTRODUCTION

Cataract is a major cause of blindness and ocular morbidity throughout the world, and any means of delaying or preventing its onset would have enormous social and economic benefits.<sup>1</sup>

Accurate biometric measurement has become a key component in surgical planning because of advancements in cataract surgery and intraocular lens (IOL) design. One of the key factors in ocular biometry is LT. It is not only a crucial variable in fourth-generation IOL formulations, but connections with other ocular factors may also have an impact on the actual position of the lens after surgery.<sup>2</sup> The ability to obtain accurate measurement of ocular biometric dimensions, such as AL, ACD, and crystalline

LT, is essential for many clinical and research applications. AL is used clinically for intraocular lens (IOL) power calculation prior to cataract and refractive surgeries and to diagnose ocular conditions such as staphyloma, to evaluate the risk of retinal detachment as well as to measure structural and dimensional components in myopia.<sup>3-6</sup>

Bearing in mind the vital role of ocular biometrics in diagnostic and therapeutic eye care, we evaluated the distribution of LT and its associations with other ocular biometric factors among cataract patients.

### METHODS

This cross-sectional study data was obtained from January 2022-June 2022 at GMC, Jammu, India. Total


978 cataract patients were retrospectively included. Ocular biometric factors were calculated by biometer and corneal analyser integrated of Topcon which included k1, k2, LT, CCT, ACD, WTW distance and AL were collected based on the medical records. The associations between LT and general or ocular factors were assessed. The inclusion criteria for this study are patients who underwent cataract surgery in the period of year January 2022-June 2022 in GMC Jammu with complete medical record data. The exclusion criteria were any co-morbidity in ophthalmology including glaucoma, retinal detachment, traumatic cataract, and any other ophthalmology condition except cataract. Medical records with missing data were excluded from the study. We analyzed the data using descriptive analysis and correlated each variable using the Spearman's Rho analysis.

**RESULTS**

Table 1 shows that amongst 978 patients, females outnumbered the males as 539 patients were females and 439 were males. Mean age of the patients was 56.36 years. In most of the cases that is 51.4% right eye was affected.

**Table 1: Demographic characteristic of the patients.**

Demographic characteristic	Value
Gender	Male 439
	Female 539
Age (Years)	Mean ± SD 56.36±15.70
Laterality (%)	Right eye 51.4
	Left eye 48.6



**Figure 1: Gender distribution.**

**Table 2: Patient's baseline characteristics.**


Baseline characteristics	Value (Mean ± SD)
LT	4.35±0.41
AL	23.17±0.73
CCT	498.01±37.23
ACD	3.28±0.33
WTW	11.41±0.45

Table 2 shows that the mean LT was 4.35 mm, AL was 23.17 mm, CCT was 498.01 mm, ACD was 3.28 mm and white to white distance was 11.41 mm.

**Table 3: Distribution of LT according to age.**

Age (Years)	Mean LT
≤30	3.68±0.38
31-40	4.21±0.42
41-50	4.29±0.40
51-60	4.34±0.63
61-70	4.62±0.65
71-80	4.68±0.44
>80	4.70±0.52

When LT was stratified by age, it gradually increased with the increasing age.




**Figure 2: Age distribution and LT.**

**Table 4: Spearman's correlation of LT with other ocular biometric factors.**

Ocular biometric factors	R	P value
AL	0.141	0.002
CCT	0.006	0.001
ACD	-0.49	0.05
WTW	0.10	0.03

On applying Spearman's correlation, a significant positive relationship of LT with AL, CCT and white to white distance was observed. This means that as there is increase in LT, AL, CCT and white to white distance will also increase. A significant negative relationship of LT with ACD was observed; this means that LT increases, ACD will decrease.



**Figure 3: The correlation between LT and other ocular parameters.**

**DISCUSSION**

Given the increasingly predictable refractive outcomes of contemporary phacoemulsification during the past two decades, surgeons have placed a growing emphasis on the use of optical biometers, which have shown to be reliable and competitive in many measurements.<sup>7,8</sup> The

availability of this technology, together with advancements in IOL platforms, has significantly increased the demand for refractive cataract surgery.<sup>9</sup>

In addition to playing an increasingly significant role in the surgical planning of cataract patients, LT is a significant biometric parameter that also correlates to morphological changes in the lens with age or cataract formation.<sup>10-13</sup>

In the present study, there were 978 eyes of preoperative cataract patients met the inclusion criteria. Amongst 978 patients, females outnumbered the males as 55.11% (539) patients were females. The mean age of the patients was 56.36±15.70 years.

In this study mean LT was 4.35±0.41mm, this is thicker than the study done by Hashemi et al where the mean LT was 4.28 mm, but when compared with the study done by Jivrajka et al it was thinner where the mean LT was 4.93±0.56 mm.<sup>14,15</sup>

Mean AL was 23.17±0.73mm, which is slightly higher than study done in the USA, South China, Iran and Myanmar where the mean AL was 23.46±1.03 mm, 23.48 mm (23.40-23.55), 23.14 mm and 22.76 mm respectively.<sup>14-17</sup>

Mean CCT was 498.01±37.23, mean ACD was 3.28±0.33 mm and white to white distance was 11.41±0.45 mm. When LT was stratified by age, it was significantly greater in older patients. The increase in LT with age, which was noted in this study, will largely explain the progressive reduction in ACD. An increase in LT with age has been observed in other studies.<sup>18,19-21</sup> Age related nuclear sclerosis, increase in protein fiber layers forming under the capsule as well as continuously lying down of lens fibers with increasing age, have been reported as being the reasons for increase in LT in cataractous lens with advancing age.<sup>22</sup>

LT was positively correlated with AL, WTW and CCT, while negatively correlated with ACD. It is possible that a bigger lens and a thicker cornea are related because the association between LT and CCT in our study is consistent with earlier results.<sup>23</sup> In addition, we found LT and ACD to be negatively correlated. The lens tends to thicken in both directions when a cataract develops, which may cause the anterior chamber to become shallower.<sup>24</sup> Despite the fact that the statistical link between the preoperative ACD and the postoperative position of the IOL has been amply demonstrated, it is possible that the IOL formula bias will exist when the preoperative ACD is used independently as opposed to in conjunction with the LT.<sup>10</sup>

The association between LT and WTW is positive, which is another intriguing finding. Due to the absence of practical measurement tools, it might be challenging to directly measure capsular bag sizes in clinical practice.

As a result, the WTW may offer useful information for these sizes. Large WTW may be linked to the growth of bigger anterior segments, indicating the incompatibility of fixed-size IOLs with large capsular bags and the potential for further IOL stability issues. As a result, the IOL should be carefully chosen in cataract patients with bigger lenses.

## CONCLUSION

LT is an important biometric parameter that should be considered, along with other biometric parameters, in determining effective lens position in patients undergoing cataract surgery. ACD is predominantly influenced by LT than the AL.

*Funding: No funding sources*

*Conflict of interest: None declared*

*Ethical approval: The study was approved by the Institutional Ethics Committee*

## REFERENCES

- Perkins ES. Lens thickness in early cataract. *Br J Ophthalmol.* 1988;72(5):348-53.
- Meng J, Wei L, He W, Qi J, Lu Y, Zhu X. Lens thickness and associated ocular biometric factors among cataract patients in Shanghai. *Eye Vis (Lond).* 2021;8(1):22.
- Rabsilber T, Jepsen C, Auffarth G, Holzer M. Intraocular lens power calculation: Clinical comparison of two ocular biometry devices. *J Cataract Refract Surg.* 2010;36:230-4.
- Saka N, Ohno-Matsui K, Shimada N, Sueyoshi S, Nagaoka N, Hayashi W et al. Long-term changes in axial length in adult eyes with pathologic myopia. *Am J Ophthalmol.* 2010;150:562-8.
- Ruiz-Moreno J, Montero J, de la Vega C, Alio J, Zapater P. Retinal detachment in myopic eyes after phakic intraocular lens implantation. *J Cataract Refract Surg.* 2006;22:247-52.
- Saw S, Chua W, Gazzard G, Goh D, Tan DTH, Stone RA. Eye growth changes in myopic children in Singapore. *Br J Ophthalmol.* 2005;89:1489-94.
- Ortiz A, Galvis V, Tello A, Viaña V, Corrales MI, Ochoa M et al. Comparison of three optical biometers: IOLMaster 500, Lenstar LS 900 and Aladdin. *In Ophthalmol.* 2019;39(8):1809-18.
- Cho YJ, Lim TH, Choi KY, Cho BJ. Comparison of ocular biometry using new swept-source optical coherence tomography-based optical biometer with other devices. *Korean J Ophthalmol.* 2018;32(4):257-64.
- Kim SY, Lee SH, Kim NR, Chin HS, Jung JW. Accuracy of intraocular lens power calculation formulas using a swept-source optical biometer. *PLoS One.* 2020;15(1):e0227638.
- Plat J, Hoa D, Mura F, Busetto T, Schneider C, Payerols A et al. Clinical and biometric determinants of actual lens position after cataract surgery. *J Cataract Refract Surg.* 2017;43(2):195-200.
- Olsen T. Prediction of the effective postoperative (intraocular lens) anterior chamber depth. *J Cataract Refract Surg.* 2006;32(3):419-24.
- Tripathi RC, Tripathi BJ. Lens morphology, aging, and cataract. *J Gerontol.* 1983;38(3):258-70.
- Klein BE, Klein R, Moss SE. Lens thickness and five-year cumulative incidence of cataracts: the Beaver Dam Eye Study. *Ophthalmic Epidemiol.* 2000;7(4):243-8.
- Hashemi H, Khabazkhoob M, Miraftab M, Emamian MH, Shariati M, Abdolahinia T et al. The distribution of axial length, anterior chamber depth, lens thickness, and vitreous chamber depth in an adult population of Shahroud, Iran. *BMC Ophthalmol.* 2012;12:50.
- Jivrajka R, Shamma MC, Boenzi T, Swearingen M, Shamma HJ. Variability of axial length, anterior chamber depth, and lens thickness in the cataractous eye. *J Cataract Refract Surg.* 2008;34(2):289-94.
- Warrier S, Wu HM, Newland HS, Muecke J, Selva D, Aung T, Casson RJ. Ocular biometry and determinants of refractive error in rural Myanmar: the Meiktila Eye Study. *Br J Ophthalmol.* 2008;92(12):1591-4.
- Chen H, Lin H, Lin Z, Chen J, Chen W. Distribution of axial length, anterior chamber depth, and corneal curvature in an aged population in South China. *BMC Ophthalmol.* 2016;16(1):47.
- Fotadar R, Wang J, Burlutsky G, Morgan I, Rose K, Wong T et al. Distribution of axial length and ocular biometry measured using partial coherence laser interferometry (IOL Master) in an older white population. *Ophthalmology.* 2010;117:417-23.
- Badmus SA, Ajaiyeoba AI, Adegbehingbe BO, Onakpoya OH, Adeoye AO. Associations between ocular biometry and anthropometric measurements in a Nigerian population. *Niger Postgrad Med J.* 2016;23:127-31.
- He M, Huang W, Zheng Y, Alsbirk PH, Foster PJ. Anterior chamber depth in elderly Chinese: The Liwan Eye Study. *Ophthalmology.* 2008;115:1286-90.
- Foster PJ, Alsbirk PH, Baasanhu J, Munkhbayar D, Uranchimeg D, Johnson GJ. Anterior chamber depth in Mongolians: Variation with age, sex, and method of measurement. *Am J Ophthalmol.* 1997;124:53-60.
- Chinawa EN, Ezech EI. The distribution of ocular biometrics among patients undergoing cataract surgery. *Niger J Ophthalmol.* 2018;26:40-5.
- Nangia V, Jonas JB, Sinha A, Matin A, Kulkarni M. Central corneal thickness and its association with ocular and general parameters in Indians: the Central India Eye and Medical Study. *Ophthalmology.* 2010;117(4):705-10.
- Jonas JB, Nangia V, Gupta R, Khare A, Sinha A, Agarwal S et al. Anterior chamber depth and its associations with ocular and general parameters in adults. *Clin Exp Ophthalmol.* 2012;40(6):550-6.

25. Zhu X, He W, Zhang Y, Chen M, Du Y, Lu Y. Inferior decentration of multifocal intraocular lenses in myopic eyes. *Am J Ophthalmol.* 2018;188:1-8.

**Cite this article as:** Sharma S, Sharma AK, Gupta S, Gupta D. Lens thickness and associated ocular biometric factors among cataract patients in tertiary hospital. *Int J Res Med Sci* 2023;11:852-6.