

## Research Article

# *In vitro* study of *Myristica fragrans* seed (Nutmeg) ethanolic extract and quercetin compound as anti-inflammatory agent

Kartika Dewi<sup>1</sup>, Budi Widyarto<sup>1</sup>, Pande Putu Erawijantari<sup>2</sup>, Wahyu Widowati<sup>1\*</sup>

<sup>1</sup>Faculty of Medicine, Maranatha Christian University, Bandung-40164, West-Java, Indonesia

<sup>2</sup>Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung-40163, West-Java, Indonesia

**Received:** 15 July 2015

**Accepted:** 11 August 2015

**\*Correspondence:**

Dr. Wahyu Widowati,

E-mail: wahyu\_w60@yahoo.com

**Copyright:** © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

### ABSTRACT

**Background:** Inflammation is one of an important biological response toward injury. Cytokine and mediator are produced by macrophage during the inflammatory process. Anti-inflammatory is important to treat the dangerous of chronic inflammation associated with chronic disease. Various plants and their derived compounds have been used in the treatment of inflammation including *Myristica fragrans*. The present study was designed to determine anti-inflammatory potential of *M. fragrans* seed (Nutmeg) ethanolic extract and pure quercetin extract from *M. fragrans* on LPS stimulated-murine macrophage cell line (RAW 264.7).

**Methods:** Cell viability assay to evaluate the non toxic concentration in cell line was performed by MTS assay. The anti-inflammatory potential was assayed through the inhibitory activity of *M. fragrans* seed extract and quercetin on NO, TNF- $\alpha$ , IL-6, and IL-1 $\beta$  production.

**Results:** The lowest cytotoxic activity and safe substance on RAW 264.7 cell were 50 and 10  $\mu\text{g/mL}$  concentration of the *M. fragrans* seed ethanolic extract and quercetin compound. *M. fragrans* dose-dependently inhibited NO, TNF- $\alpha$ , IL-6 and IL-1 $\beta$  production on LPS stimulated-RAW 264.7. The 50  $\mu\text{g/mL}$  of *M. fragrans* seed ethanolic extract showed the highest TNF- $\alpha$ , IL-6, IL-1 $\beta$  and nitrite-associated with NO inhibitory activity.

**Conclusions:** This research suggested that *M. fragrans* seed extract and quercetin compound possess the anti-inflammatory potential showed through the inhibition of TNF- $\alpha$ , IL-6, IL-1 $\beta$  and NO secretion.

**Keywords:** Anti-inflammatory, *Myristica fragrans*, Quercetin, Cytokine, RAW 264.7

### INTRODUCTION

Inflammation is the complex biological response to overcome the harmful stimuli which aimed as host defense and usually activated in most disease conditions including pathogen exposure, cell injury and irritation.<sup>1,2</sup> Inflammation relates to various diseases such as rheumatoid arthritis, inflammatory bowel disease, atherosclerosis, Alzheimer's, and has a role in various kinds of cancer.<sup>3</sup> Macrophage plays an important role in Reactive Oxygen Species (ROS), Reactive Nitrogen Species (RNS), cytokines [Interleukine (IL)-1 $\beta$ , IL-6, Tumor Necrosis Factor (TNF)- $\alpha$ ] production and Nitric

Oxide (NO) mediated inflammation and prostaglandin. Chronic inflammation is a dysregulated response to persistent noxious stimuli and seems to be related to tissue malfunction. Prolonged inflammation is associated with some chronic human disorders including cancer, allergy, arthritis, atherosclerosis and autoimmune disease, so that anti-inflammatory agent become important.<sup>4,5</sup>

Various substances from plants are playing beneficial role in the prevention and even treatment of different diseases.<sup>6,7</sup> Plants extract are rich of bioactive chemicals and most of them free from adverse effects.<sup>8</sup> Some

herbals shows anti-inflammatory potential.<sup>9</sup> Lipopolysaccharide (LPS)-stimulated-murine macrophage cell line (RAW 264.7) is an appropriate model for evaluating and screening of anti-inflammatory agents from plants extract.<sup>10</sup> Macrophages in mammalian immune system play a significant role in immediate defense against foreign agents.<sup>11</sup> LPS is a component of the cell walls of gram negative bacteria that the most powerful activators of macrophages and involves the production of pro-inflammatory cytokines.<sup>12</sup>

*M. fragrans* is one of medicinal plant that have variety of active phytochemical including vitamins, alkaloids, flavonoids, lignans and phenolic, etc. These compounds render their effects via different mechanism such as metal chelation, inhibition of lipid peroxidation and quenching of singlet oxygen to act as antioxidants.<sup>13</sup> The utility of *Myristica fragrans* has been known used as spice in Indonesia and introduced to Europe. *M. fragrans* has aromatic, stimulant, narcotic, carminative, astringent, aphrodisiac, hypolipidaemic, antithrombotic, anti-platelet aggregation, antifungal, antidysenteric, and anti-inflammatory activities.<sup>14</sup> Quercetin is one of the major classes of phytochemicals in *M. fragrans* seeds. Quercetin, the most commonly occurring flavonoid is an excellent antioxidant that is also suggested to possess other beneficial activities.<sup>15,16</sup> The aim of this research is to determine anti-inflammatory potential of *M. fragrans* seed (Nutmeg) ethanolic extract and pure quercetin extract from *M. fragrans* on LPS stimulated-murine macrophage cell line (RAW 264.7).

## METHODS

### Plant extract preparation

This study was performed to continue the previous study of the antioxidant properties of spices extracts including *M. fragrans* seed extract.<sup>17</sup> The dried seeds of *M. fragrans* were crushed then extracted using ethanol 70%. Extraction was performed using maceration technique with ethanol 70% as a solvent. Every 24 hours the ethanol filtrate was filtered and collected until the filtrate become colorless. Evaporation was performed using 40°C evaporator until the pasta form product was obtained. The extracts were store at 4°C.<sup>18,19</sup> The *M. fragrans* extract and quercetin compound from *M. fragrans* extract was used as the treatment.

### Cell culture

The murine macrophage cell line RAW 264.7 (ATCC®TIB-71™) was given by Biomolecular and Biomedical Research Center, Aretha Medika Utama. The RAW 264.7 cells were grown and maintained in DMEM (Dulbecco's Modified Eagle Medium) supplemented with 20% FBS, 100 U/mL penicillin (Invitrogen), 100 µg/mL streptomycin (Invitrogen). The culture was incubated at 37°C in humidified atmosphere and 5% CO<sub>2</sub> until the cells were confluent. The cells then washed and

resuspended in DMEM. The cells then seeded on plates and treated using *M. fragrans* extract in different concentration (100 mg/mL, 50 mg/mL, 25 mg/mL, 0 mg/mL) and different concentration of quercetin (100 µM, 50 µM, and 25 µM).<sup>20</sup>

### Viability assay

MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium) assay (Promega, Madison, WI, USA) is a colorimetric method for determining the number of viable cells in proliferation or cytotoxicity assays.<sup>21</sup> The cells were seeded in 96-well plate (5 x 10<sup>3</sup> cells per well) in 100 µL medium (DMEM supplemented with 10% FBS and 100 U/mL penicillin and streptomycin) and incubated for 24 hours at 37°C in a humidified atmosphere and 5% CO<sub>2</sub>. The medium then washed and supplemented with 99 µL new medium and 1 µL of *M. fragrans* extract in various concentration (100 mg/mL, 50 mg/mL, 25 mg/mL, 0 mg/mL), quercetin in various concentrations (100 µM, 50 µM, and 25 µM) and DMSO in different plate then incubated for 24 hours. Untreated cells served as the control. 20 µL MTS was added to each well. The plate was incubated in 5% CO<sub>2</sub> at 37°C incubator for 4 hours. The absorbance was measured at 490 nm on microplate reader. The data are presented as the percentage of viable cells (%).<sup>18,19</sup> The viability assay was performed to determine the suitable concentration for the next assay.

### Pro-inflammatory activation of cells

Inflammation cell triggered using modified method from Yoon et al.<sup>10</sup> and Khan et al.<sup>22</sup> The cells were seeded in 6-well plate (5 x 10<sup>3</sup> cells per well) and incubated for 24 hours at 37°C in a humidified atmosphere and 5% CO<sub>2</sub>. The medium (DMEM supplemented with 10% FBS and 100 U/mL penicillin and streptomycin) then washed and supplemented with 1600 µL growth medium and 200 µL extract solution or quercetin in different concentration then incubated for 1-2 hours. 200 µL LPS (1 µg/mL) was added into the medium and incubated for 24 hours at 37°C in a humidified atmosphere and 5% CO<sub>2</sub>. The growth medium was taken for assay and centrifuged at 2000 g for 10 minutes. Supernatant was stored at -79°C.

### Quantification of nitrite concentration and inhibitory activity assay

Abnova kit (No cat. KA 1342) was used to determine the concentration of nitrite associated with NO production. After pre-incubation of RAW 264.7 cells with LPS and *M. fragrans* extract/quercetin for 24 h, the quantity of nitrite accumulated in the culture medium (DMEM supplemented with 10% FBS and 100 U/mL penicillin and streptomycin) was measured as an indicator of NO production. 200 µL assay buffer was added into the blank well and 100 µL of standard solution with 100 µL assay buffer was added into standard. Briefly, 100 µL of cell

medium was mixed with 100  $\mu$ L assay buffer. The mixture was incubated at room temperature for 10 minutes and the absorbance at 540 nm was measured in a microplate reader. The quantity of nitrite was determined from sodium nitrite standard curve. The LPS-stimulated cells without extract or quercetin was used as positive control. The normal cell was used as negative control.

#### **Quantification of TNF- $\alpha$ concentration and inhibitory activity assay**

BioLegend kit (No. cat 421701) was used to determine the TNF- $\alpha$  concentration in the culture medium.<sup>23</sup> The plate was washed four times using 300  $\mu$ L wash buffer then 50  $\mu$ L matrix C was added into the standard well. 50  $\mu$ L assay buffer was added into each of the sample well. 50  $\mu$ L sample was added into each of the sample well and 50  $\mu$ L standard solution was used as standard. The LPS-stimulated cells without *M. fragrans* extract or quercetin was used as positive control. The normal cell was used as negative control. The plate then wrapped and incubated for 2 hours in 200 rpm orbital shaker. All of the solution was washed by 200  $\mu$ L wash buffer, after two hours of incubation. 100  $\mu$ L Rat-TNF- $\alpha$  detection antibody was added to each well and incubated in 100 rpm orbital shaker. The plate then washed again using 200  $\mu$ L wash buffer, after one hour of incubation. 100  $\mu$ L avidin HRP was added in each well and incubated for 30 minutes in 200 rpm orbital shaker. The plate then washed five times using 200  $\mu$ L wash buffer. 100  $\mu$ L substrate solution F was added into each well and incubated in the dark condition for 10 minutes. Stop solution was added into each well and the results then readed using MultiSkan Go ThermoScientific ELISA reader in the 450 nm wavelength.

#### **Quantification of IL-6 concentration and inhibitory activity assay**

The amount of IL-6 in the culture supernatant was measured with a commercial kit (mouse max standard set, BioLegend). Plate was washed using 300  $\mu$ L wash buffer four times, 50  $\mu$ L matrixes C was added in to standard well and 50  $\mu$ L assay buffer was added into sample well. 50  $\mu$ L samples were introduced in sample well and 50  $\mu$ L standard solutions were added in standard well then incubated in orbital shaker 200 ppm for 2 hour at room temperature. The LPS-stimulated cells without *M. fragrans* extract or quercetin was used as positive control. The normal cell was used as negative control. 100  $\mu$ L of Rat IL-6 detection antibody was added then incubated for 1 hours in room temperature 200 rpm orbital shaker. The solution then discharged and plate was washed using 200  $\mu$ L of wash buffer four times. 100  $\mu$ L avidin HRP solution was added and kept at room temperature for 30 minutes in orbital shaker 200 ppm. Plate was washed again for 5 times and 100  $\mu$ L substrate solution F was added then kept for 10 min in dark room. After 10 minutes, 100  $\mu$ L stop solution was added and

absorbance was read by using Miniskan ELISA Reader at 450 nm.<sup>24</sup>

#### **Quantification of IL-1 $\beta$ concentration and inhibitory activity assay**

The IL-1 $\beta$  concentration was determined based on BioLegend (No.Cat 432601) kit protocol. 300  $\mu$ L wash buffer was used to wash the plate for times before the addition of 50  $\mu$ L matrix C into standard well and 50  $\mu$ L of assay buffer into sample well. 50  $\mu$ L standard solution was added into standard well and 50  $\mu$ L sample was added into the sample well. The LPS-stimulated cells without *M. fragrans* extract or quercetin was used as positive control. The normal cell was used as negative control. Plate then incubated in room temperature 100 rpm orbital shaker. After two hours of incubation the solution was discharged and washed four times using 200  $\mu$ L wash buffer. 100  $\mu$ L of substrat solution F then added to the each well then incubated in dark condition. Stop solution the added into each well to terminite the reaction after 10 minutes of incubation. The absorbance was readed using this MultiSkan Go ThermoScientific ELISA reader in 450 nm of wavelength.<sup>25</sup>

#### **Statistical analysis**

All data were derived from three independent experiments. Statistical analysis was conducted using SPSS software (version 17.0). Value were presented as mean  $\pm$  SD. Significant differences between the groups were determined using the analysis of variat (ANOVA) and Tukey Post Hoc test. Statistical significance was set at P<0.05.

## **RESULTS**

#### **Effects of *M. fragrans* on RAW 264.7 cells viability**

RAW 264.7 cell viability assay was the preliminary study to test the effect of *M. fragrans* seed extract and quercetin compound toward RAW 264.7 cell viability. The aims of this assay is to determine the safe and non toxic concentration for the next assay. Viability was measured by MTS assay based on the conversion of yellow tetrazolium salt to form a purple formazan product.<sup>21</sup> The percent of cells viability was determined by comparing the cells viability value of treatments to the control. This viability assays (Table 1) shows that *M. fragrans* seed extract and quercetin in given concentration still available for the normal RAW 264.7 cells except for the concentration of 100  $\mu$ g/mL. In the highest concentration the RAW 264.7 cells viability was low, indicate that concentration is little bit toxic to the cells so that concentration will not used in the next step of assay.

### ***M. fragrans* seed extract exhibited the higher nitric oxide (NO) inhibitory activity**

The positive control of this test shows the highest concentration of nitrite concentration compared to the negative control and extract or quercetin treated cells (Table 2). That indicate the LPS successfully induce the inflammation on the RAW 264.7 cells.<sup>26</sup> The percent of inhibitory activity was determine by the value of positive

control nitrite concentration minus the nitrite concentration of treatment divided to the nitrite concentration of positive control. 50 µg/mL *M. fragrans* seed ethanolic extracts shows the highest nitrite inhibitory activity (Table 2). The negative control shows the highest value of nitrite inhibitory too because the normal cell was used. The normal cell did not release much nitrite like LPS-stimulated cells so the inhibitory activity become high.

**Table 1: Mean and Tukey HSD post hoc test of RAW 264.7 cell's viability of various concentration extract and quercetin measured in triplicate.**

| Samples                         | Viability                 |                           |                           |
|---------------------------------|---------------------------|---------------------------|---------------------------|
|                                 | 100 µg/mL                 | 50 µg/mL                  | 10 µg/mL                  |
| <i>M. fragrans</i> seed extract | 85.27 ± 1.34 <sup>a</sup> | 90.84 ± 2.08 <sup>b</sup> | 97.75 ± 1.34 <sup>c</sup> |
| Quercetin                       | 83.21 ± 6.4 <sup>a</sup>  | 92.34 ± 5.78 <sup>b</sup> | 99.52 ± 9.47 <sup>c</sup> |

Data were presented as mean ± standard deviation. Different superscript letters (<sup>a,b,c</sup>) in the same row (concentrations) are significant at P<0.05 (Tukey HSD post hoc test)

**Table 2: Mean and Tukey HSD post hoc test of nitrite content and nitrite inhibitory activity over the positive control of various concentration extract and quercetin measured in triplicate.**

| Samples                                  | Nitrite oxide              |                                 |
|--|----------------------------|---------------------------------|
|  | Nitrite concentration (µM) | Nitrite inhibitory activity (%) |
| Positive control                         | 35.68 ± 0.23 <sup>f</sup>  | 0.0067 ± 0.65 <sup>a</sup>      |
| Negative control                         | 4.92 ± 0.49 <sup>a</sup>   | 86.2250 ± 1.37 <sup>f</sup>     |
| <i>M. fragrans</i> seed extract 50 µg/mL | 21.27 ± 0.03 <sup>b</sup>  | 40.3883 ± 0.091 <sup>e</sup>    |
| <i>M. fragrans</i> seed extract 10 µg/mL | 25.38 ± 0.10 <sup>c</sup>  | 28.8733 ± 0.29 <sup>d</sup>     |
| Quercetin 50 µg/mL                       | 25.81 ± 0.10 <sup>d</sup>  | 27.6650 ± 0.27 <sup>c</sup>     |
| Quercetin 10 µg/mL                       | 30.84 ± 0.14 <sup>e</sup>  | 13.5733 ± 0.38 <sup>b</sup>     |

Data were presented as mean ± standard deviation. Different superscript letters (<sup>a,b,c,d,e,f</sup>) in the same row (nitrite concentration and nitrite inhibitory activity) are significant at P<0.05 (Tukey HSD post hoc test)

### ***M. fragrans* seed extract posses the higher TNF-α inhibitory activity**

Tumor necrosis factor-α (TNF-α) plays an important role as the main cytokine of inflammation processes. The highest TNF-α concentration was shown by the positive control. The *M. fragrans* ethanolic seed extract or quercetin-treated cells shows the lower TNF-α concentration compared to the positive (Table 3). The

percent of inhibition was determine by using the value of positive control TNF-α concentration minus the TNF-α concentration of treated cells compare to the TNF-α concentration of positive control. The *M. fragrans* seed extract in 50 µg/mL concentration have the highest inhibitory activity (Table 3). The negative control-LPS-nonstimulated cells show the high value of TNF-α inhibitory activity because it did not release much TNF-α like the LPS-stimulated cells so the inhibition value become high.

**Table 3: Mean and Tukey HSD post hoc test of TNF-α concentration and TNF-α inhibitory activity over the positive control of various concentration extract and quercetin measured in triplicate.**

| Samples                                  | TNF-α                       |                               |
|--|-----------------------------|-------------------------------|
|  | TNF-α concentration (pg/mL) | TNF-α inhibitory activity (%) |
| Positive control                         | 500.17 ± 7.74 <sup>f</sup>  | 0.00 ± 1.55 <sup>a</sup>      |
| Negative control                         | 251.20 ± 14.06 <sup>a</sup> | 49.78 ± 2.81 <sup>f</sup>     |
| <i>M. fragrans</i> seed extract 50 µg/mL | 280.90 ± 6.20 <sup>b</sup>  | 43.84 ± 1.24 <sup>e</sup>     |
| <i>M. fragrans</i> seed extract 10 µg/mL | 348.20 ± 10.36 <sup>c</sup> | 30.38 ± 2.07 <sup>d</sup>     |
| Quercetin 50 µg/mL                       | 368.30 ± 15.92 <sup>d</sup> | 26.37 ± 3.18 <sup>c</sup>     |
| Quercetin 10 µg/mL                       | 438.73 ± 13.83 <sup>e</sup> | 12.28 ± 2.77 <sup>b</sup>     |

Data were presented as mean  $\pm$  standard deviation. Different letters (a,b,c,d,e,f) in the same row (TNF- $\alpha$  concentration and inhibitory activity) are significant at  $P < 0.05$  (Tukey HSD post hoc test)

#### Both of *M. fragrans* seed extract and quercetin possess IL-6 inhibitory activity

*M. fragrans* seed extract and quercetin possess the IL-6 inhibitory activity in concentration dependent manner. The highest inhibitory activity was showed by *M. fragrans* seed extract in the concentration of 50  $\mu\text{g/mL}$

(Table 4). IL-6 is one of cytokine that takes a part in acute inflammation including hematopoiesis, 'response immune' regulation, and inflammation.<sup>24,27</sup> The negative control-LPS-nonstimulated cells show the high value of IL-6 inhibitory activity because it did not release much IL-6 like the LPS-stimulated cells so the inhibition value become high when it compared to the positive control.

**Table 4: Mean and Tukey HSD post hoc test of IL-6 concentration and IL-6 inhibitory activity over the positive control of various concentration extract and quercetin measured in triplicate.**

| Samples   | IL-6                              |                                |
|---|-----------------------------------|--------------------------------|
|   | IL-6 concentration (pg/mL)        | IL-6 inhibitory activity (%)   |
| Positive control                                    | 912.14 $\pm$ 201.31 <sup>b</sup>  | 0.00 $\pm$ 0.00 <sup>a</sup>   |
| Negative control                                    | 110.81 $\pm$ 12.93 <sup>ab</sup>  | 87.34 $\pm$ 3.55 <sup>c</sup>  |
| <i>M. fragrans</i> seed extract 50 $\mu\text{g/mL}$ | 44.76 $\pm$ 10.04 <sup>a</sup>    | 94.83 $\pm$ 2.03 <sup>c</sup>  |
| <i>M. fragrans</i> seed extract 10 $\mu\text{g/mL}$ | 693.47 $\pm$ 176.32 <sup>ab</sup> | 24.32 $\pm$ 2.57 <sup>ab</sup> |
| Quercetin 50 $\mu\text{g/mL}$                       | 307.10 $\pm$ 17.13 <sup>ab</sup>  | 65.32 $\pm$ 7.10 <sup>bc</sup> |
| Quercetin 10 $\mu\text{g/mL}$                       | 890.90 $\pm$ 409.59 <sup>ab</sup> | 9.73 $\pm$ 51.65 <sup>ab</sup> |

Data were presented as mean  $\pm$  standard deviation. Different letters (a,b,ab,bc) in the same row (IL-6 concentration and inhibitory activity) are significant at  $P < 0.05$  (Tukey HSD post hoc test)

#### Both of *M. fragrans* seed extract and quercetin exhibited IL-1 $\beta$ inhibitory activity

IL-1 $\beta$  expression as pro-inflammatory cytokine is enhanced following crush injury, infection, invasion and

become an important sign of inflammation.<sup>28</sup> *M. fragrans* seed extract possess the highest activity in inhibiting IL-1 $\beta$ . The negative control-LPS-nonstimulated cells show the high value of IL-1 $\beta$  inhibitory activity because it did not release IL-1 $\beta$  much like the LPS-stimulated cells so the inhibition value become high (Table 5).

**Table 5: Mean and Tukey HSD post hoc test of IL-1 $\beta$  concentration and IL-1 $\beta$  inhibitory activity over the positive control of various concentration extract and quercetin measured in triplicate.**

| Samples   | IL-1 $\beta$                       |                                      |
|---|------------------------------------|--------------------------------------|
|   | IL-1 $\beta$ concentration (pg/mL) | IL-1 $\beta$ inhibitory activity (%) |
| Positive control                                    | 51.93 $\pm$ 8.60 <sup>a</sup>      | 0.00 $\pm$ 0.00 <sup>a</sup>         |
| Negative control                                    | 80.77 $\pm$ 23.83 <sup>a</sup>     | 54.38 $\pm$ 30.88 <sup>ab</sup>      |
| <i>M. fragrans</i> seed extract 50 $\mu\text{g/mL}$ | 59.97 $\pm$ 16.78 <sup>a</sup>     | 100.10 $\pm$ 98.25 <sup>b</sup>      |
| <i>M. fragrans</i> seed extract 10 $\mu\text{g/mL}$ | 112.36 $\pm$ 43.60 <sup>a</sup>    | 14.56 $\pm$ 18.67 <sup>ab</sup>      |
| Quercetin 50 $\mu\text{g/mL}$                       | 74.97 $\pm$ 18.31 <sup>a</sup>     | 43.66 $\pm$ 20.01 <sup>ab</sup>      |
| Quercetin 10 $\mu\text{g/mL}$                       | 72.84 $\pm$ 28.63 <sup>a</sup>     | 10.00 $\pm$ 5.26 <sup>ab</sup>       |

Data were presented as mean  $\pm$  standard deviation. Different letters (a,b,ab) in the same row (IL-1 $\beta$  concentration and inhibitory activity) are significant at  $P < 0.05$  (Tukey HSD post hoc test)

## DISCUSSION

Inflammation is one of an important biological response toward injury. Cytokine and mediator are produced by macrophage during the inflammatory process.<sup>29</sup> Anti-inflammatory is important to treat the dangerous of chronic inflammation associated with chronic disease. Natural phytochemicals plays a significant role in drug discovery. Various plants and their derived compounds

have been used in the treatment of inflammation.<sup>9</sup> Anti-inflammatory drugs should have effect on prime causative factors, inhibitory effect or blocking effect on initial reaction set in a biological model by the prime cause and thereby inhibit the established inflammation, and effect on end results of established inflammation.<sup>30</sup> In this *in vitro* study, we evaluated the anti-inflammatory potential of *Myristica fragrans* seed (nutmeg) extract and the quercetin compound from *M. fragrans* through NO,

TNF- $\alpha$ , IL-6 and IL-1 $\beta$  inhibitory activity assays in LPS stimulated-murine macrophages cell line (RAW 264.7). Lipopolysaccharide (LPS) is a pro-inflammatory glycolipid component of the gram negative bacterial cell wall that stimulated inflammation process.<sup>31</sup> The activation of macrophage by LPS leads to secretion of inflammatory molecules such as the inflammatory cytokine including TNF- $\alpha$ , IL-6, IL-1 $\beta$  and the free radical NO, which play an important role in inflammation.<sup>32,33</sup>

The *M. fragrans* seed extract and quercetin compound in 10 and 50  $\mu\text{g/mL}$  of concentration displayed positive result and no toxicity to RAW 264.7. Non toxicity of that substrate was indicated by high percentage of viable cells in viability test. Viability test is an important aspect of pharmacology that deals with the adverse effect of bioactive substance on living organism prior to the use as drug or chemical in clinical use.<sup>34-36</sup> The *M. fragrans* seed extract and quercetin compound from *M. fragrans* dose-dependently inhibited NO, TNF- $\alpha$ , IL-6 and IL-1 $\beta$  production on LPS stimulated-RAW 264.7.

Tumor Necrosis Factor (TNF- $\alpha$ ) is one of the proinflammatory cytokines that plays an important role in inflammation.<sup>37</sup> TNF- $\alpha$  is involved in inflammatory response by activating Nuclear Factor kappa-light-chain-enhancer of activated B cells (NF- $\kappa\text{B}$ ), cytokine, and adhesion molecule inducer.<sup>5,38,39</sup> Increased concentration of TNF- $\alpha$  are believed to cause the cardinal signs of inflammation to occur. TNF- $\alpha$  will stimulate white cell phagocytosis and the production of inflammatory lipid prostaglandin E2 (PGE2).<sup>40</sup> The TNF- $\alpha$  inhibitory activity measurement is important in anti-inflammatory potential agent screening since this cytokine is an important mediator of inflammation.<sup>31</sup> Blocking one cytokine can be sufficient as anti-inflammatory because cytokine exist in cascades.<sup>37</sup> In particular, TNF- $\alpha$  may stimulate additional inflammatory pathways result in additional inflammatory pathways resulting Nitric Oxide (NO).<sup>5</sup> IL-6 act as both pro- and anti-inflammatory cytokines.<sup>41,42</sup> IL-6 along with TNF- $\alpha$  and IL-1 $\beta$  serve as endogenous pyrogens that causes fever during inflammation by up regulating the inflammatory responses and stimulating the production of acute phase reactants.<sup>43</sup>

*M. fragrans* seed ethanolic extract and quercetin compound from *M. fragrans* showed the inhibitory activity potential on TNF- $\alpha$ , IL-6 and IL-1 $\beta$  production. Our study is in the line with study showing that macelignan as an active compound of *M. fragrans* can targeted NF- $\kappa\text{B}$  and COX-2 gene. NF- $\kappa\text{B}$  induced inflammation by activating several cytokine, chemokine and the gene including TNF- $\alpha$ , IL-1 $\beta$ , IL-6, IL-8, COX-2, 5-LOX and iNOS.<sup>44-46</sup> Inhibition on NF- $\kappa\text{B}$  could inhibit the production of TNF- $\alpha$ , IL-6, and IL-1 $\beta$  because there is positive feedback between NF- $\kappa\text{B}$  and that cytokine.<sup>47</sup> Recent research also confirmed that quercetin as one of active compound from *M. fragrans* can inhibit various cytokines including TNF- $\alpha$ , IL-6, and IL-1 $\beta$ .<sup>15,16,31,48,49</sup>

NO plays a significant role in host immune defense, vascular regulation, neurotransmission, and other system in normal condition. Overproduction of inducible NO Synthase (iNOS) is especially related to various human diseases including inflammation.<sup>50</sup> LPS can induce iNOS transcription and transduction with subsequent NO production in murine macrophage RAW 264.7 cells. Among the oversecreted inflammatory mediator, NO has been strongly implicated the pathogenesis of several disease.<sup>51</sup> NO inhibitions activity is important for anti-inflammatory agent screening. Our result showed that *M. fragrans* seed ethanolic extract and quercetin compound from *M. fragrans* can inhibit the nitrite production-an indicator of nitric oxide (NO) synthesis. Cao et al. (2013) identified six compounds (licarin B, 3'-methoxylicarin B, myrisfrageal A, isodihydrocainatidin, dehydrodiisoeugenol, and myrisfrageal B) of *M. fragrans* that showed inhibition of nitric oxide production in LPS-stimulated RAW 264.7.<sup>52</sup> Our result also in line with the research that showed quercetin can inhibit NO production and iNOS protein and mRNA expression.<sup>53</sup> *M. fragrans* seed contains several active phytochemicals including isoeugenol, methyl-eugenol, eugenol, dihydroguaiaretic acid,  $\gamma$ -terpinene, terpinen-4-ol, myristic acid, oleanolic acid, palmitic acid, camphene, lauric acid, myrecne, kaempferol, and also quercetin as the most common compound found in the *M. fragrans* seed.<sup>54</sup>

## CONCLUSION

This research suggested that *M. fragrans* seed extract and quercetin compound possess the anti-inflammatory potential showed by the inhibition of TNF- $\alpha$ , IL-6, IL-1 $\beta$  and NO production. It is well known that NO and PGE<sub>2</sub> are main macrophage-derived inflammatory mediators.<sup>50,55,56</sup> Furthermore, it has been reported that TNF- $\alpha$ , IL-6 and IL-1 $\beta$  is an important inflammatory cytokine. Therefore, it has been thought suppression on TNF- $\alpha$ , IL-6, IL-1 $\beta$  and NO is a good strategy to reduce inflammation.<sup>50</sup> Validation of active samples in animal models should eventually follow for further study.

## ACKNOWLEDGEMENTS

We gratefully acknowledge the financial support of the Research Center and Service Community, Maranatha Christian University, Bandung, Indonesia for research grant 2014. This study was successfully done by the facilities support and design study by Maesaroh Maesaroh and Nurul Fauziah from Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung, West Java, Indonesia.

*Funding:* The study was funded by Research grant 2014, Research Center and Service Community, Maranatha Christian University

*Conflict of interest:* None declared

*Ethical approval:* Not required

## REFERENCES

1. Phanse M, Patil M, Abbulu K, Chaundhari P, Patel B. *In vivo* and *in vitro* screening of medicinal plants for their anti-inflammatory activity: an overview. JAPS. 2012;2(6):19-33.
2. Joseph J, Bindhu A, Aleykutty N. *In vitro* and *in vivo* antiinflammatory activity of *Clerodendrum paniculatum* Linn Leaves. Indian J Pharm Sci. 2013;75(3):376-9.
3. Fang S, Hsu C, Yen G. Anti-inflammatory effects of phenolic compound isolated from the fruits of *Artocarpus heterophyllus*. J Agr Food Chem. 2008;56(12):4463-8.
4. Medzhitov R. Origin and physiological roles of inflammation. Nature. 2008;454:428-35.
5. da Silveira e Sa R, Andrade L, de Oliveira RdRB, de Sousa D. A review on anti-inflammatory activity of phenylpropanoids found in essential oils. Molecules. 2014;19:1459-80.
6. Luximon R, Baborun T, Crozier A. Antioxidant actions and phenolic and vitamin C contents of common Mauritian exotic fruits. J Sci Food Agric. 2003;83:496-502.
7. Hanna L, Maria L, Jerzy D, Ratiporn H, Sumitra P, Yong-Seo P, et al. Bioactive properties of snake fruit (*Salacca edulis* Reinw) and mangosteen (*Garcinia mangostana*) and their influence on plasma lipid profile and antioxidant activity in rats feed cholesterol. Eur Food Res Technol. 2006;223:697-703.
8. Lee H, Kim S, Lee C, Ahn Y. Cytotoxic and mutagenic effects of *Cinnamom cassia* bark derived materials. J Microbiol Biotechnol. 2004;14(6):1176-81.
9. Agnihotri S, Wakode S, Agnihotri A. An overview on anti-inflammatory properties and chemo-profiles of plants used in traditional medicine. Indian J Nat Prod Resour. 2010 Jun;1(2):150-67.
10. Yoon WJ, Ham Y, Kim SS, Yoo BS, Moon JY, Baik J, et al. Suppression of pro-inflammatory cytokines, iNOS and COX-2 expression by brown algae *Sargassum micracanthum* in RAW 264.7 macrophages. Eur Asia J Biol Sci. 2009;3:130-43.
11. Joo T, Sowndhararajan K, Hong S, Lee J, Park S, Kim S, et al. Inhibition of nitric oxide production in LPS-stimulated RAW 264.7 cells by stem bark of *Ulmus pumila* L. Saudi J Biol Sci. 2014;21:427-35.
12. Nicholas C, Batra S, Vargo M, Voss O, Gavrilin M, Wewers M, et al. Apigenin blocks lipopolysaccharide-induced lethality *in vivo* and proinflammatory cytokines expression by inactivating NF-kappaB through the suppression of p65 phosphorylation. J Immunol. 2007;179:7121-7.
13. Gupta A, Bansal V, Babu V, Maithil N. Chemistry, antioxidant and antimicrobial potential of nutmeg (*Myristica fragrans* Houtt). JGEB. 2013;11:25-31.
14. Muchtaridi, Subarnas A, Apriyanto A, Mustarichie R. Identification of compounds in the essential oil of nutmeg seeds (*Myristica fragrans* Houtt) that inhibit locomotor activity in mice. Int J Mol Sci. 2010;11:4771-81.
15. Nair M, Mahajan S, Reynolds J, Aalinkeel R, Nair H, Schwartz S, et al. The flavonoids quercetin inhibits proinflammatory cytokine (tumor necrosis factor alpha) gene expression in normal peripheral blood mononuclear cells via modulation of NF-kappabeta system. Clin Vaccine Immunol. 2006;13:319-28.
16. Manjeet K, Ghosh B. Quercetin inhibits LPS-induced nitric oxide and tumor necrosis factor-alpha production in murine macrophages. Int J Immunopharmacol. 1999;21:435-43.
17. Widowati W, Ratnawati H, Husin W, Maesaroh M. Antioxidant properties of spice extracts. In: International Conference on Advance Molecular Bioscience and Biomedical Engineering. Malang, East Java: ICAMBBE; 2014.
18. Widowati W, Wijaya L, Wargasetia T, Bachtiar I, Yellianty Y, Laksmiawati D. Antioxidant, anticancer and apoptosis-inducing effects of Piper extracts in HeLa cells. J ExpIntegr Med. 2013;3(3):225-30.
19. Widowati W, Mozef T, Risdian C, Yellianty Y. Anticancer and free radical scavenging potency of *Catharanthus roseus*, *Dendrophtho epetandra*, *Piper bettle* and *Curcuma mangga* extracts in breast cancer cell line. Oxidant Antiooxid. 2013;2(2):137-42.
20. Lee J, Park W. Anti-inflammatory effect of myristicin on RAW 264.7 macrophages stimulated with polyinosinic-polycytidylic acid. Molecules. 2011;16:7132-42.
21. Malich G, Markovic B, Winder C. The sensitivity and specificity of the MTS tetrazolium assay for detecting the *in vitro* cytotoxicity of 20 chemicals using human cell lines. Toxicology. 1997;124:179-92.
22. Khan T, Wagener J, Bost T, Martinez J, Accurso F, Riches D. Early pulmonary inflammation in infants with cystic fibrosis. AJRCCM. 1995;151(4):1075-82.
23. Kilanisa K, Maizels N. Ultrasensitive isolation, identification, and quantification of DNA-protein adducts by ELISA-based RADAR assay. Nucleic Acids Res. 2014;108:1-12.
24. Kuroishi T, Bando K, Endo Y, Sugawara S. Metal allergens induce nitric oxide production by mouse dermal fibroblast via the hypoxia-inducible factor-2alpha-dependent pathway. Toxicol Sci. 2013;135(1):119-28.
25. Muarayam M, Kakuta S, Maruhashi T, Shimizu K, Seno A, Kubo S, et al. CTRP3 plays an important role in the development of collagen-induced arthritis in mice. Biochem Biophys Res Commun. 2014;443(1):42-8.
26. Sarkar D, Fisher P. Molecular mechanism of aging-associated inflammation. Cancer Lett. 2006;236:13-23.
27. Ishikara K, Hirano T. IL-6 in autoimmune disease and chronic inflammatory proliferative disease. Cytokine Growth Factor Rev. 2002;13:357-68.

28. Zhang J, An J. Cytokines, inflammation, and pain. *Int Anesthesiol Clin*. 2007;45(2):27-37.
29. Jung C, Jung H, Shin Y, Park J, Jun C. *Eleutherococcus senticosus* extract attenuates LPS-induced iNOS expression through the inhibition of Akt and JNK pathways in murine macrophage. *J Ethnopharmacol*. 2007;113:183-7.
30. Naik S, Sheth U. Inflammatory process and screening methods for anti-inflammatory agents - a review. *J Postgrad Med*. 1976;22:5-21.
31. Boots A, Wilms L, Swennen E, Kleinjans J, Bast A, Haenen G. *In vitro* and *ex vivo* anti-inflammatory activity of quercetin in healthy volunteers. *Nutrition*. 2008;24(7-8):703-10.
32. Gunawardena D, Shanmugam K, Govindaraghavan S, Low M, Bennet L, Head R, et al. Determination of anti-inflammatory activities of standardised preparation of plant and mushroom-based foods. *Eur J Clin Nutr*. 2014;53(1):335-43.
33. Soromou L, Zhang Z, Li R, Chen N, Guo W, Huo M, et al. Regulation of inflammatory cytokines in lipopolysaccharide-stimulated RAW 264.7 murine macrophage by 7-O-methyl-naringenin. *Molecules*. 2012;17:3574-85.
34. Rajalakshmi A, Jayachitra A, Gopal P, Krithiga N. Toxicity analysis of different medicinal plant extracts in Swiss albino mice. *Biomed Res*. 2014;1(2):1-6.
35. Subramanian L, Zuraini Z, Yen C, Yee L, Lachimanan Y, Screenivasan S. Acute oral toxicity of methanolic seed extract of *Cassia fistula* in mice. *Molecules*. 2011;16:5268-82.
36. Lalitha P, Shubashini K, Jayanthi P. Acute toxicity of extracts of *Eichhornia crassipes* (MART.) SOLMS. *Asian J Pharm Clin Res*. 2012;5(4):59-61.
37. Dinarello C. Anti-inflammatory agents: present and future. *Cell*. 2010;140(6):935-50.
38. Tak P, Firestein G. NF-kappaB: a key role in inflammatory disease. *J Clin Invest*. 2001;107:7-11.
39. Libby P. Inflammation in atherosclerosis. *Nature*. 2002;420:868-74.
40. Maroon J, Bost J, Maroon A. Natural anti-inflammatory agents for pain relief. *Surg Neuro Int*. 2010;1:80.
41. Jones S, Horiuchi S, Topley N, Yamamoto N, Fuller G. The soluble interleukin 6 receptor: mechanisms of production and implications in disease. *FASEB J*. 2001;15:43-58.
42. Wong P, Campbell I, Egan P, Ernst M, Wicks I. The role of interleukin-6 family of cytokines in inflammatory arthritis and bone turnover. *Arthritis Rheum*. 2003;48(5):1177-89.
43. Damte D, Reza M, Lee S, Jo W, Park S. Anti-inflammatory activity of dichloromethane extract of *Auricularia auricula-judae* in RAW264.7 cells. *Toxicol Res*. 2011;27(1):11-4.
44. Aggrawal B, Prasad S, Reuter S, Kannappan R, Yadav V, Park B, et al. Identification of novel anti-inflammatory agents from ayurvedic medicine for prevention of chronic diseases. *Curr Drug Targets*. 2011;12(11):1595-653.
45. Ghosh S, May E. NF-B and Rel proteins: evolutionarily conserved mediators of immune responses. *Annu Rev Immunol*. 1998;16:225.
46. Navdeep S, Wendy C, David S, Paul T. Role of oxidant in NF-kappaB activation and TNF-alpha gene transcription induced by hypoxia and endotoxin. *J Immunol*. 2000;165(2):1013-21.
47. Kagoya Y, Yoshimi A, Kataoka K, Nakagawa M, Kumano K, Arai S, et al. Positive feedback between NF-kappa B and TNF-alpha promotes leukemia-initiating cell capacity. *JCI*. 2014;124(2):528-42.
48. Mueller M, Hobiger S, Jungbauer A. Anti-inflammatory activity of extracts from fruits, herbs and spices. *Food Chem*. 2010;122:987-96.
49. Bellik Y, Boukraa L, Alzahrani H, Bakhotmah B, Abdellah F, Hammoudi S, et al. Molecular mechanism underlying anti-inflammatory and anti-allergy activities of phytochemicals: an update. *Molecules*. 2013;18:322-53.
50. Kang CH, Choi Y, Choi I, Lee J, Kim GY. Inhibition of Lipopolysaccharide-Induced iNOS, COX-2, and TNF-alpha expression by aqueous extract of *Orixa japonica* in RAW 264.7 cells via suppression of NF-kappaB activity. *Trop J Pharm Res*. 2011;10(2):161-8.
51. MacMicking J, Xie Q, Nathan C. Nitric oxide and macrophage function. *Ann Rev Immunol*. 1997;15:323-50.
52. Cao G, Yang X, Xu W, Li F. New inhibitors of nitric oxide production from the seeds of *Myristica fragrans*. *Food Chem Toxicol*. 2013;62:167-71.
53. Hamalainen M, Nieminen R, Vuorela P, Heinonen M, Moilanen E. Anti-inflammatory effects of flavonoids: genistein, kaempferol, quercetin, and daidzein inhibit STAT-1 and NF-KB Activations, whereas flavone, isohamnetin, naringenin, and pelargonidin inhibit only NF-KB activation. *Mediators Inflamm*. 2007;2007:45673.
54. Suhaj M. Spice antioxidants isolation and their antiradical activity: a review. *J Food Compos Anal*. 2006;19:531-7.
55. Vodovotz Y, Kim P, Ermentrout G, Chow C, Bahar I, Billiar T. Inflammatory modulation of hepatocyte apoptosis by nitric oxide: *in vivo*, *in vitro*, and *in silico* studies. *Curr Mol Med*. 2004;4:753-62.
56. Moncada S, Higgs E. Nitric oxide and the vascular endothelium. *Handb Exp Pharmacol*. 2006;176:213-54.

**Cite this article as:** Dewi K, Widyarto B, Erawijantari PP, Widowati W. *In vitro* study of *Myristica fragrans* seed (Nutmeg) ethanolic extract and quercetin compound as anti-inflammatory agent. *Int J Res Med Sci* 2015;3:2303-10.