

Research Article

Serum gamma glutamyl transferase levels in association with lipids and lipoproteins in type2 diabetes mellitus

Duggirala Rajarajeswari*, T. Sharmila Krishna, M. Prasad Naidu, Jupalli Nagayya Naidu

Department of Biochemistry, Narayana Medical College, Nellore-524003, Andhra Pradesh, India

Received: 21 March 2014

Revised: 13 May 2014

Accepted: 23 May 2014

*Correspondence:

Dr. Duggirala Rajarajeswari,

E-mail: bio.rajeswari@gmail.com

© 2014 Rajarajeswari D et al. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Serum gamma glutamyl transferase is widely used as a marker for alcohol induced liver disease. Recently it has gained importance due to its role in type2 diabetes mellitus. A raised serum gamma glutamyl transferase level indicates hepatic steatosis and visceral fat deposition, leading to insulin resistance and diabetes. In the present study we examined the association between serum gamma glutamyl transferase levels with lipids and lipoproteins in diabetes mellitus.

Methods: The study was carried out on 50 subjects with type 2 diabetes mellitus and compared with 50 age and sex matched healthy controls attending outpatient department of general medicine, Narayana medical college, Nellore. Serum gamma glutamyl transferase was measured by calorimetric kinetic assay. Fasting blood sugar was measured by glucose oxidase method using automated analyzer. Serum triglycerides, total cholesterol and high density lipoprotein are measured by standard enzymatic procedures and low density lipoprotein by Friedwald equation.

Results: Serum gamma glutamyl transferase levels in diabetic cases were significantly elevated compared to normal healthy controls ($P < 0.001$). There was a positive correlation between gamma glutamyl transferase, lipids and low density lipoprotein and inverse correlation with high density lipoprotein ($r = -0.298$).

Conclusion: Our results suggest a possible role of gamma glutamyl transferase in the pathophysiology and progression of type 2 diabetes mellitus.

Keywords: Serum gamma glutamyl transferase, lipids, Blood glucose, Type 2 diabetes mellitus

INTRODUCTION

There is a growing body of evidence that serum Gamma Glutamyl Transferase (GGT) is a predictor of Type 2 Diabetes Mellitus (T2DM) independent of factors associated with excess alcohol consumption and liver disease.¹ Serum GGT is a cell surface protein and plays a key role in glutathione metabolism. Glutathione is the main intracellular thiol antioxidant agent in mammalian cells.² Serum GGT activity is affected by both genetic and environmental factors.³ Various tissues produce GGT which differ in their sugar moieties. Most of the serum GGT is derived from liver tissue and binds to carriers like

lipoproteins and albumin.^{4,5} A number of recent studies have suggested that abnormal hepatocellular function is associated with obesity, insulin resistance, and type 2 diabetes.^{6,7} The loss of a direct effect of insulin to suppress hepatic glucose production and glycogenolysis in the liver causes an increase in hepatic glucose production.⁸ Raised liver enzymes reflect chronic ectopic fat deposition.⁹ Serum GGT may be a simple and reliable marker of visceral and hepatic fat deposition and hepatic steatosis which can lead to hepatic insulin resistance and long term hepatic insulin resistance may lead to T2DM.¹⁰ In this context the study was under taken to study the serum GGT levels and its possible associations with

triglycerides, total cholesterol, Low Density Lipoprotein (LDL) and High Density Lipoprotein (HDL) in T2DM.

METHODS

The study was carried out on 50 subjects with T2DM and 50 age and sex matched healthy controls recruited from general medicine out-patient department, of Narayana medical college and hospital, Nellore, A.P. All the subjects underwent a complete medical evaluation by a physician. Informed and written consent was obtained from all subjects. Both male and female subjects between 18-65 years of age were included in the study. Pregnant, lactating women, alcoholics, patients with liver disease and patients on insulin and other drugs which might influence lipid and glucose levels and liver enzymes were excluded from the study.

Biochemical parameters

Fasting blood sugar is measured by glucose oxidase method using automated chemistry (Humaster-300, GMBH, and Germany) analyzer using Human kits. Serum triglycerides, total cholesterol and HDL cholesterol are analyzed by standard enzymatic procedures and LDL cholesterol by Friedwald equation. Serum GGT is measured by calorimetric kinetic assay.

Statistical analysis is performed using SPSS software (16.0). P value <0.05 is considered as statistically significant.

RESULTS

All subjects are grouped into normal and diabetic cases. Table 1: Demonstrates the mean \pm SD values of the main characteristics of diabetic cases and controls. Serum GGT was significantly elevated in diabetic cases (Mean \pm SD 67.20 ± 22.97) compared to normal healthy controls (Mean \pm SD 24.71 ± 7.91) P value <0.001. Subjects with diabetes had significantly elevated levels of fasting blood sugar, total cholesterol, triglycerides and LDL cholesterol but HDL cholesterol is significantly decreased in diabetics compared to healthy controls. Correlation between serum GGT and biochemical characteristics in subjects with and without diabetes: The analysis for correlation of serum GGT with biochemical characteristics of the subjects indicated a positive correlation between GGT and lipids and LDL lipoprotein but inverse correlation to HDL lipoprotein ($r = -0.298$). Figure 1 demonstrates the correlation between serum GGT and fasting blood sugar. There is a significant positive correlation between serum GGT and fasting blood sugar in subjects with diabetes ($r = 0.35$) Figure 2. Shows a strong positive correlation between serum GGT and triglycerides in subjects with T2DM ($r = 0.112$).

Figure 3 shows a strong negative association between serum GGT and HDL in subjects with T2DM ($r = -0.298$).

Table 1: Main characteristics of diabetic cases and controls.

Biochemical parameters	Cases n=50	Controls n=50	P value
GGT (IU/L)	67.20 ± 22.97	24.71 ± 7.91	0.0001
Fasting blood sugar (mg/dl)	160.9 ± 33.23	88.88 ± 6.27	0.0001
Cholesterol (mg/dl)	408.68 ± 84.53	157.02 ± 27.18	0.0001
Triacyl glycerol (mg/dl)	247.76 ± 127.25	85.78 ± 31.23	0.0001
HDL(mg/dl)	30.42 ± 5.24	79.94 ± 15.35	0.0001
LDL(mg/dl)	156 ± 12.52	93.72 ± 19.99	0.0001

Data are expressed as mean+ standard deviation. P <0.05 were significant




Figure 1: Association between serum GGT and fasting blood glucose in subjects with T2DM.




Figure 2: Association between serum GGT and triglycerides in subjects with T2DM.




Figure 3: Association between serum GGT and HDL lipoprotein in subjects with diabetes.

DISCUSSION

GGT has been known mainly as a marker of alcohol consumption and liver diseases.¹¹ Serum GGT within normal ranges has been reported to be associated with risk factors of cardiovascular diseases and components of metabolic syndrome.¹² Many studies hypothesized to link GGT and diabetes.¹³ This study was done to estimate serum GGT levels and analyze its association with blood glucose, lipids and lipoproteins in subjects with diabetes and compared with age and sex matched healthy controls. Studies in humans and animal models have indicated that serum GGT levels were significantly elevated in diabetes.¹⁴ In our study we observed that serum GGT levels were significantly elevated in subjects with diabetes compared to normal controls ($P < 0.0001$). Similar studies on the relationship between liver enzymes and diabetes in both sexes in general population have found higher levels of serum GGT in diabetes. Our results demonstrated that increased serum GGT levels were associated with increased levels of glucose in diabetes. Recent studies on diabetic middle aged men and women have shown increased levels of GGT when compared with liver enzymes, alcohol consumption, and body mass index. Serum GGT was also correlated with insulin resistance-markers, waist-circumference, triglycerides, fasting plasma glucose, HbA_{1c}, systolic and diastolic blood pressure. In parallel to this our study also observed association of serum GGT with blood glucose, lipids and lipoproteins; however in our study we have excluded subjects with a history of alcoholism. Our results are consistent with those of previous studies and indicate that elevated serum GGT is associated with an increased risk of diabetes. One explanation for our findings is that the elevation of serum GGT could be an expression of excess fat deposited in liver, which is regarded as a feature of the insulin resistance syndrome. We observed a significant correlation between GGT, lipids, lipoproteins and blood sugar levels. In our study slightly positive association between serum GGT and fasting blood sugar ($r = 0.08$) was observed. Previous study on patients with diabetes after a three year follow-up period had showed that raised GGT was correlated with central obesity, increased fasting glucose, triglycerides, and blood pressure in both sexes. In subjects with T2DM a strong positive association between serum GGT and triglycerides ($r = 0.112$), serum GGT and LDL lipoprotein ($r = 0.05$), serum GGT and cholesterol ($r = 0.027$) was observed. We also observed a strong negative association between serum GGT and HDL lipoprotein ($r = -0.298$). Association of GGT with lipids and lipoproteins could be explained by the antioxidant property of GGT.^{15,16} Elevated GGT could reflect subclinical inflammation, which represents the underlying oxidative stress. GGT levels are closely related to oxidative stress because cellular GGT has a central role in glutathione homeostasis by initiating the breakdown of extracellular glutathione, a critical antioxidant defense for the cell.^{17,18} Our study shows that serum GGT levels are significantly associated with

T2DM, serum triglycerides, total cholesterol and LDL lipoprotein and inversely correlated to HDL lipoprotein accelerating the progression of T2DM related cardiovascular risk. Our study high lightens the role of serum GGT in developing T2DM and can be used as a marker to diagnose T2DM.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the institutional ethics committee

REFERENCES

1. André P, Balkau B, Charles MA, Eschwège E. Gamma-glutamyl transferase activity and development of the metabolic syndrome, in middle-aged men and women: the DESIR cohort. *Diabet Care*. 2007;30(9):2355-61.
2. Whitfield JB. Gamma glutamyl transferase. *Crit Rev Clin Lab Sci*. 2000;138(4):263-355.
3. Benini F, Pigozzi MG, Baisini O, Romanini L, Ahmed H, Pozzi A, et al. Increased serum gamma-glutamyl-transpeptidase concentration is associated with non-alcoholic steatosis and not with cholestasis in patients with chronic hepatitis C. *J Gastroenterol Hepatol*. 2007;22(10):1621-26.
4. Huseby NE. Hydrophilic form of γ -glutamyl transferase: proteolytic formation in liver homogenates and its estimation in serum. *Clin Chim Acta*. 1982;124(1):113-21.
5. Grundy, Scott M. Gamma-glutamyl transferase another biomarker for metabolic syndrome and cardiovascular risk. *Arterioscl Thromb Vasc Biol*. 2007;27(1):4-7.
6. Sabanayagam C, Shankar A, Li J, Pollard, Ducatman A. Serum gamma-glutamyl transferase level and diabetes mellitus among US adults. *Eur J Epidemiol*. 2009;24(7):369-73.
7. Lim JS, Lee DH, Park JY, Jin, Jacobs. A strong interaction between serum γ -glutamyl transferase and obesity on the risk of prevalent type 2 diabetes: results from the third national health and nutrition examination survey. *Clin Chem*. 2007;53(6):1092-8.
8. Duckworth, William C, Frederick G. Hamel, Daniel E. Peavy. Hepatic metabolism of insulin. *Am J Med*. 1988;85.5:71-6.
9. Kasapoglu B, Turkay C, Bayram Y, Koca C. Role of GGT in diagnosis of metabolic syndrome: a clinic-based cross-sectional survey. *Indian J Med Res*. 2010;132(1):56-61.
10. Lippi G, Targher G, Montagnana M, Salvagno GL, Guidi GC. Relationship between gamma-glutamyl transferase, lipids and lipoprotein(a) in the general population. *Clin Chim Acta*. 2007;384:163-6.
11. Vozarova B, Stefen N, Lindsey RS, Saremi A, Pratley RE, Bogardus C, et al. High alanine aminotransferase is associated with decreased hepatic insulin sensitivity and predicts the

- development of type 2 diabetes. *Diabetes.* 2002;51:1889-95.
12. Vartiainen E, Jousilahti P, Alfthan G, Sundvall J, Pietinen P, Puska P. Cardiovascular risk factor changes in Finland 1972-1997. *Int J Epidemiol.* 200;29:49-56.
 13. Koro CE, Bowlin SJ, Bourgeois N, Fedder DO. Glycemic control from 1988 to 2000 among US adults diagnosed with type 2 diabetes a preliminary report. *Diabet Care.* 2004;27(1):17-20.
 14. Lee Duk-Hee, Myron D. Gross, David R. Jacobs. Association of serum carotenoids and tocopherols with γ -glutamyltransferase: the cardiovascular risk development in young adults (CARDIA) Study. *Clin Chem.* 2004;50.3:582-8.
 15. Duk-Hee Lee, Rune Blomhoff, David R. Jacobs Jr. Is serum gamma glutamyl transferase a marker of oxidative stress? *Free Radic Res.* 2004;38.6:535-9.
 16. Sabanayagam C, Shankar A, Li J, Pollard C, Ducatman A. Serum gamma-glutamyl transferase level and diabetes mellitus among US adults. *Eur J Epidemiol.* 2009;24:369-73.
 17. Iqbal Azhar, Uzma Iftikhar, Farah Amir Ali, Shakoor Memon, Nudrat Zuberi. Comparison of gamma glutamyl transferase in normal and in type 2 diabetics. *J Pak Med Assoc.* 2010;60(11):945-8.
 18. Giovanni Targher, Lorenz Bertolini, Felice Poli, Stefano Rodella, Luca Scala, Tessari R, et al. Non-alcoholic fatty liver disease and risk of future cardiovascular events among type 2 diabetic patients. *Diabetes.* 2005;54(12):3541-6.

DOI: 10.5455/2320-6012.ijrms20140819

Cite this article as: Rajarajeswari D, Sharmila Krishna T, Prasad Naidu M, Naidu JN. Serum gamma glutamyl transferase levels in association with lipids and lipoproteins in type2 diabetes mellitus. *Int J Res Med Sci* 2014;2:838-41.