

## Original Research Article

# Implication incidence age groups in cancer patients who suffer from death rumination disorder and employed psychological defense styles

Minoo Sharbafshaaer<sup>1\*</sup>, Mohammad A. Mashhadi<sup>2</sup>

<sup>1</sup>Department of Psychology, University of Sistan and Baluchestan, Zahedan, Sistan and Baluchestan, Iran

<sup>2</sup>Department of Haematology and Oncology, Zahedan University of Medical Sciences, Zahedan, Sistan and Baluchestan, Iran

**Received:** 02 October 2016

**Accepted:** 26 November 2016

### \*Correspondence:

Dr. Minoo Sharbafshaaer,

E-mail: [minoshaer@gmail.com](mailto:minoshaer@gmail.com)

**Copyright:** © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

### ABSTRACT

**Background:** Cancer is malignant disease with high risk of incidences which disease lead to stressful and pressure condition. This study was done to analyse age groups in cancer patients who suffer from death rumination disorder and employed psychological defense styles.

**Methods:** The study was done using two questionnaires such as: death rumination questionnaire and defense style questionnaire-60 (DSQ-60). Study was conducted in 200 patients who after oncologist diagnosis bedded in hospital. Investigation used all data collected analysed by SPSS version 21 software, the MANOVA test was used.

**Results:** In particular, 46-55 years old was in high risk of age group in cancer population, there was a statistical significant difference on death rumination disorder and employing psychological defense styles with regard to age. Age groups correlated with death focus factor of death rumination disorder. Age rank correlated with sublimation, self-assertion, splitting-self, help-rejecting, undoing, affiliation factors of defense styles.

**Conclusions:** Findings of this study revealed that there is difference between age groups, death rumination disorder and psychological defense styles in cancer patients. Death rumination is response of distress and death thinking which make by cancer diseases, then defense styles in patients encounter for adaptive with pressure and stressful condition of diseases.

**Keywords:** Age groups, Cancer patients, Death rumination, Psychological defense styles

### INTRODUCTION

Nowadays cancer disease has attracted worldwide attention in all age rank. Cancer is spread abnormal cells, primary and next serious effect is tumor and metastatic of malignant cells.<sup>1</sup> According to the IRNA, it estimated cancer disease was second cause of death in Iran.<sup>2</sup> Patients with cancer diagnosis ought to give chemical therapy for treatment. Owing to the fact, situation and condition of treatment follow by high stress and death rumination. Therefore patients using psychological defense mechanisms to adaptive with disease condition and treatment meanwhile they respond stress. Actually,

defense mechanisms happen in every patient.<sup>3</sup> In this study, we aim to look into estimated difference age groups in term of cancer patient's scores on death rumination disorder and employed psychological defense styles.

### METHODS

The present investigation was conducted in 200 patients that they went into hospital and they bedded in oncology parts of hospital after diagnosis with cancer disease. Cases of cancer study getting various treatments such as surgery, chemotherapy and radiology. Patients should be

informed and consent to answered interview and questionnaire. Interview of each patient took approximately 45-60 minutes to complete the questionnaires. The study started from 12<sup>th</sup> April 2014 to 20<sup>th</sup> March 2015. Data collection was during 12 months.

**Instrument**

Two questionnaires were used including death rumination questionnaire and defense style questionnaire. Death rumination questionnaire was on the basis of Ruminative Responses Scale. RRS made by Nolon-Hoeksema. Death rumination questionnaire semi structured interviews and self-report inventories, baseline negative mood and self-criticism, this involving repetitive self-focused.<sup>4,5</sup>

It estimates 6 version of death rumination include: self-focus, self-critical, self-exhaust, death-focus, death-analysis, and dysfunctional-mood. Respondents answer each of the 13 items on a 5 point Likert scale with click of one (not at all applicable to me) and five (completely applicable to me). Defense style questionnaire (DSQ-60) created by Thygesen, Drapeau, Trijsburg, Lecours and de Roten, 2008. Self-report is cornerstone of DSQ questionnaire, defense style based in the 60 item model of the scale that it assesses 30 psychological defense styles such as: acting-out, affiliation, altruism, anticipation, denial, devaluation of self, devaluation of other, displacement, dissociation, fantasy, help-rejecting complaining, humour, idealization, intellectualization, isolation, omnipotence, passive aggressive, projection, identification, rationalization, reaction formation, repression, self-assertion, self-observation, splitting of self, splitting of other, sublimation, suppression, undoing, and withdrawal. Respondents answer each of the 60 items on a 9 point Likert scale with anchors of one (not at all applicable to me) and nine (completely applicable to me).<sup>6</sup>

The All questionnaires were checked for completion of information and the responses were coded for entry in the computer. Questionnaire data analyse was performed with Microsoft the statistical package for the social science (SPSS-pc) software version 21. Age rank was considering with frequency statistics and multivariate test assessment age rank with 30 styles of defense and 6 death rumination kinds, furthermore, all statistical tests were done to determine any significant difference at 5% level.

**RESULTS**

Total study patients were 200, in six age group from 18 till 83 years old, they bedded in hospital. study evaluate in cancer patients, they got various treatment meanwhile patients suffer from 6 type death rumination disorder hence they employed 30 kind of psychological defense mechanisms (mentioned in methodology part) consciously and unconsciously. Described of age groups in cancer patients research was 100 per cent. Age mean in cancer patients (44.74) years old and SD (16.95) with

95% interval confidence. Evidence indicate 46-55 age group was peak of incidence age groups which get more attention in cancer patients’ population, the 36-45 years old was next vulnerability age group (Table 1).

**Table 1: Incidence cancer in age groups.**

Group	Variable (years old)	Frequency	Percent
1	18-25	31	15.5
2	26-35	34	17
3	36-45	42	21
4	46-55	44	22
5	56-70	31	15
6	71-83	10	9
Total		200	100

(95% confidence interval)

Multivariate analysis was found that there was a statistically significant difference between six rank of age groups cancers’ patients with death rumination disorder and employing psychological defense style,  $F(180.749)=1.42$ ,  $p=0.001$ ; Wilks’ Lambda=0.25; partial eta squared=0.24, Bonforroni adjusted alpha level= 0.001 (Table 2).

**Table 2: Effect age difference in death rumination and employed psychological defense styles.**

Wilks’ Lambda value square	F	P value	Partial Eta
0.25	1.423	0.001	0.242

(99% confidence interval)

Studies shows age groups related to number factors of death rumination disorder and psychological defense styles.

Factors related to age groups including: death focus was factor of death rumination disorder and sublimation, self-assertion, splitting-self, help-rejecting, undoing and affiliation were factors of psychological defense style. Self-assertion and affiliation were 99% confidence interval. Death-focus, sublimation, splitting –self, help-rejecting and undoing was 95% confidence interval (Table 3).

**Table 3: Age groups relation with factors of death rumination and defense styles.**

Factors	F	P value	Partial Eta square
Death-focus	2.93	0.014	0.07
Sublimation	2.28	0.048	0.05
Self-assertion	3.22	0.008	0.07
Splitting-self	2.38	0.04	0.06
Help-rejecting	2.46	0.035	0.06
Undoing	2.43	0.037	0.06
Affiliation	5.67	0.001	0.12

## DISCUSSION

Present researches indicate 46 to 55 years old was major incidence age in cancer. According to the IRNA, it estimated cancer disease was second cause of death in Iran.<sup>2</sup> American Cancer Society Inc in 2015 reported, "cancer most commonly develops in older people; 78% of all cancer diagnoses are in people 55 years of age or older".<sup>7</sup> Researcher in 2015 shows, "cancer occurs in adults aged and younger than 50 years, however, cancer is high risk for people".<sup>8</sup>

National Cancer Registry, in 2015 found, "the median age of cancer patients at diagnosis will increase for all invasive cancers combined the percentage of patients aged over 70 is expected to increase, for females, from 31% in 2015 to 44% in 2025 and, for males, from 40% to 51%".<sup>9</sup> Liver cancer report in 2015 as a result indicated, "there will not only be more cancer patients, but they will be older on average. cancer in people living in less developed countries in Asia and Africa compared with those in more developed countries worldwide, the disease can develop at a younger age (typically around the age of 40)".<sup>10</sup>

Study indicates kind cancers was effective impact in death rumination and employing psychological defense styles for cancer patients, hence, death rumination disorder as high as kind cancer which is in high incidence rate. Death rumination is self-reflection of death thinking. There is highlight distinction between rumination and more adaptive forms of self-reflection, based in cognitive deficits or rumination, then, this correlated with neural and genetic state meanwhile, self-reflection could be power intervention against rumination feeling.<sup>11</sup> Rumination could be defined as prolong process and inflexible cognitive style.<sup>12</sup> Patients with negative feeling lead to death rumination especially death focus in treatment situation. There are synergistic effects of rumination and negative emotion progressively propagate and magnify then these predicting number of impulsive behaviors subsequently reported.<sup>13</sup>

Death rumination is hopeless response in patients, meanwhile death rumination related with part of brain and mental health. Rumination is typical of pathological grief responses in disaster condition.<sup>14</sup> Distinguishing of rumination condition and the other mental states is connectivity changes involving the amygdala part, it is to be important found.<sup>15</sup> Rumination is worse functioning on all mental health, this make Sleep problems.<sup>16</sup> Actually death rumination is responses of anxiety and depression mood to help patient to handle dissonance state and problem solving. Other studies implicate rumination as a mechanism of stress sensitivity and suggest pathways through which it may maintain depression and anxiety in everyday life.<sup>17</sup>

Oncologist diagnosis leads to stressful and depression mood hence patient engage death rumination. Stressful

life events induce rumination, this is potentially useful targets that is preventing the onset of depression and anxiety.<sup>18</sup> Study shows the mediational effects of adaptive and maladaptive rumination in the relationship between illness perception and negative emotions.<sup>19</sup> Patients with oncologist diagnosis whereas suffer from death rumination disorder, patients persuade to using psychological defense styles due to patients adaptive with problem in conflict condition. Psychological defense styles improved by following dissonance experience because they lead to learn professional capacity and knowledge against error event.<sup>20</sup>

Developmental, personality, and social psychologists implication from defense styles which define psychological functioning.<sup>21</sup> Therefore, mechanisms defined as unconscious process, cognitive operations alter by developmental periods for protective function, and that can be assessment of personality and experimental schedules.<sup>22</sup> Individuals can promotion behavior in certain situation, patients need to adaptive with treatment condition.<sup>23</sup>

Hemodialysis patients use defense styles as psychological treatment against stressor and improve health quality of life then defense styles amenable patients for disease treatment.<sup>24</sup> Patients with hypertension diagnosis they widely encounter from defense styles and more developed ability for feeling.<sup>25</sup> Intestinal stoma patients indicate physical and psychological health problem, patients reflected level of self-image adaptation, meanwhile elderly patients use only a small part of defense styles as coping process.<sup>26</sup>

There is strong correlation between somatic and psychological data was appeared, as part of an original complex psychosomatic model ,patients with oncologist diagnosis using defense styles as coping strategies and emotional response to all condition of diagnosis.<sup>27</sup> end-stage renal patients employing passive-aggressive behaviors which is from defense style, this related to patients personality.<sup>28</sup> There is clinical attention toward ego defense mechanisms as indicators of distress and lowered survival in cancer patients, that the maturity of adaptive mechanisms must be controlled in behavioral-treatment of cancer patients.<sup>29</sup> Defense mechanisms were general acceptable regulation in patient populations hence, clinicians can use this as most effectively target defenses in psychotherapy.<sup>30</sup> Actually, defense styles happen in every patient.<sup>3</sup>

## CONCLUSION

Findings of this study revealed that there is difference between age groups, death rumination disorder and psychological defense styles in cancer patients. Death rumination is response of distress and death thinking which make by cancer diseases, then defense styles in patients encounter for adaptive with pressure and stressful condition of diseases.

## ACKNOWLEDGEMENTS

Authors would like to thank all staffs and patients of Ali Ebne Abitaleb Hospital of their kind cooperation in their research.

*Funding: No funding sources*

*Conflict of interest: None declared*

*Ethical approval: The study was approved by the Institutional Ethics Committee*

## REFERENCES

- Adamo M, Dickie, L, Ruhl J. First course of therapy. In: National cancer institute, editor. SEER program coding and staging manual. 2015. 1<sup>st</sup> ed. Bethesda: Effective with cases diagnosed; 2015:119.
- Abadan IR. Cancer is second cause of death. 2014. Available at <http://www.irna.ir/fa/News/81291147>. Accessed 16 July 2016.
- Taylor JB. Psychological adaptive mechanisms: ego defense recognition in Practice and Research. *Psychosomatics*. 2014;55:210-1.
- Kasch KL, Klein DN, Lara ME. Construct validation study of the Response Styles Questionnaire Rumination scale in participants with a recent-onset major depressive episode. *Psychological Assessment*. 2001;13:375-83.
- Broderick PC. Mindfulness and Coping with Dysphoric Mood: Contrasts with Rumination and Distraction. *Cog Thera Rese*. 2005;29:501-10.
- Thygesen KL, Drapeau M, Trijsburg RW, Lecours S, Roten YD. Assessing defense styles: factor structure and psychometric Assessing defense styles: factor structure and psychometric properties of the new defense style questionnaire 60 (DSQ-60). *Int J Psycho Psycho Ther*. 2008;8:171-81.
- Cancer Facts and Figures 2015. Atlanta: American Cancer Society. 2015. Available at <http://www.cancer.org/research/cancerfactsstatistics/cancerfactsfigures2015/>.
- Siegel RL, Miller DK, Ahmedin J. Cancer Statistics 2015. *Ca cancer J Clin*. 2015;65:5-29.
- Cancer projections for Ireland 2015-2040. Cork: National Cancer Registry, 2014. Available at <http://www.ncri.ie/publications/cancer-trends-and-projections/cancer-projections-ireland-2015-%E2%80%932040/>. Accessed 16 July 2015.
- Diet, Nutrition, Physical Activity and Liver Cancer 2015. London: World Cancer Research Fund International's Continuous Update Project (CUP) 2015. Available at <http://www.wcrf.org/int/research-we-fund/continuous-update-project-findings-reports/liver-cancer/>.
- Nolen-Hoeksema S, Wisco BE, and Lyubomirsky S. Rethinking Rumination. *Perspect Psychol Sci* 2008;3:400-24.
- Davis RN, Nolen-Hoeksema S. Cognitive inflexibility among ruminators and nonruminators. *Cogni Ther Rese*. 2000;24:699-711.
- Selby EA, Kranzler A, Panza E, Fehling KB. Bidirectional-Compounding effects of rumination and negative emotion in predicting impulsive behavior: implications for emotional cascades. *J Pers*. 2014.
- Papa A, Rummel C, Garrison-Diehn C, Sewell MT. Behavioral activation for pathological grief. *Death Studies*. 2013;37:913-36.
- Milazzo AC, Ng B, Jiang H, Shirer W, Varoquaux G, Poline JB, et al. Identification of mood-relevant brain connections using a continuous, subject-driven rumination paradigm. *Cereb Cortex*. 2014;24:255.
- Borders A, Rothman DJ, McAndrew LM. Sleep problems may mediate associations between rumination and PTSD and depressive symptoms among OIF/OEF veterans. *Psychological Trauma: Theory, Research, Practice, and Policy*. 2015;7:76-84.
- Ruscio AM, Gentes EL, Jones JD, Hallion LS, Coleman ES, Swendsen J. Rumination predicts heightened responding to stressful life events in major depressive disorder and generalized anxiety disorder. *J Abnor Psycho*. 2015;124:17-26.
- Michl LC, McLaughlin KA, Shepherd K, Nolen-Hoeksema S. Rumination as a mechanism linking stressful life events to symptoms of depression and anxiety: longitudinal evidence in early adolescents and adults. *Abnormal Psychology*. 2013;122:339-52.
- Lu Y, Tang C, Liow CS, Ng WW, Ho CS, Ho RC. Regression analysis of maladaptive rumination, illness perception and negative emotional outcomes in Asian patients suffering from depressive disorder. *Asian J Psych*. 2014;13:1-15.
- Laurent A, Aubert L, Chahraoui KH, Bioy A, Mariage A, Quenot JP. Error in intensive care: psychological repercussions and defense mechanisms among health professionals. *Critical Care Medicine*. 2014;42:2370-8.
- Cramer P. Defense mechanisms in psychology today: Further processes for adaptation. *American Psychol*. 2000;55:637-46.
- Cramer P. Defense Mechanisms: 40 Years of Empirical Research. *Journal of Personality Assessment*. 2014;57:114-22.
- Rezki M, Bangun YR. Create the EQ modelling instrument based on Goleman and Bar-on models and psychological defense mechanisms. *Social and Behavioral Sciences*. 2014;115:394-406.
- Carvalho AF, Ramirez SP, Macêdo DS, Sales PM, Rebouças JC, Daher EF, et al. The psychological defensive profile of hemodialysis patients and its relationship to health-related quality of life. *J Nerv Ment Dis*. 2013;201:621-8.
- Pervichko E, Zinchenko Y. Ego defense mechanisms in patients with "hypertension at work" and patients with essential hypertension: a comparative analysis. *Nervous and Mental Disease*. 2013;201:621-8.

26. Ortiz-Rivas MK, Moreno-Pérez NE, Vega-Macías HD, Jiménez-González Mde J Navarro-Elías Mde G. Adaptation of self-image level and defense mechanisms in elderly patients with complicated stoma. *Enferm Clin J.* 2014;24:339-44.
27. Beresford TP, Alfors J, Mangum L, Clapp L, Martin B. Cancer survival probability as a function of ego defense (adaptive) mechanisms versus depressive symptoms. *Psychosomatics.* 2006;47:247-53.
28. Hyphantis T, Katsoudas S, Vouclari S. Ego mechanisms of defense are associated with patients' preference of treatment modality independent of psychological distress in end-stage renal disease. *Patient Prefer Adherence.* 2010;4:25-32.
29. Stepanchuk E, Zhirkov A, Yakovleva A. The coping strategies, psychological defense mechanisms and emotional response to the disease in Russian patients with chronic leukemia. *Social and Behavioral Sciences.* 2013;86:248-55.
30. Olson TR, Perry JC, Janzen JJ, Petraglia J, Presniak MD. Addressing and interpreting defense mechanisms in psychotherapy: general considerations. *Psychiatry.* 2011;74:142-65.

**Cite this article as:** Sharbafshaaer M, Mashhadi MA. Implication incidence age groups in cancer patients who suffer from death rumination disorder and employed psychological defense styles. *Int J Res Med Sci* 2016;4:5372-6.